

# Dualizing Projected Model Counting

Sibylle Möhle and Armin Biere

Institute for Formal Models and Verification  
Johannes Kepler University Linz

AVM / RiSE Workshop 2018  
September 2018



$x$	$y$	$F(x, y)$
0	0	0
0	1	1
1	0	1
1	1	1

$x$	$y$	$F(x, y)$	$\exists y.F(x, y)$
-----	-----	-----------	---------------------

$x$	$y$	$F(x, y)$	$\exists y.F(x, y)$
0	0	0	1
0	1	1	
1	0	1	1
1	1	1	

# Projected Model Counting Generalized

$F(X, Y)$  (arbitrary) propositional formula over variables  $X$  and  $Y$  with  $X \cap Y = \emptyset$

$X$  *relevant* input variables

$Y$  *irrelevant* input variables

# Projected Model Counting Generalized

$F(X, Y)$  (arbitrary) propositional formula over variables  $X$  and  $Y$  with  $X \cap Y = \emptyset$

$X$  *relevant* input variables

$Y$  *irrelevant* input variables

We are interested in the number of models projected onto  $X$ :

$$\# \exists Y. F(X, Y)$$

# Projected Model Counting Generalized

$F(X, Y)$  (arbitrary) propositional formula over variables  $X$  and  $Y$  with  $X \cap Y = \emptyset$

$X$  *relevant* input variables

$Y$  *irrelevant* input variables

We are interested in the number of models projected onto  $X$ :

$$\#\exists Y. F(X, Y)$$

Example  $F(X, Y) = x \vee y$

$$X = \{x\}$$

$$Y = \{y\}$$

$$\mathcal{M}(\exists Y. F(X, Y)) = \{x, \neg x\}$$

$$\#\exists Y. F(X, Y) = 2$$

# Projected Model Counting Generalized

$F(X, Y)$  (arbitrary) propositional formula over variables  $X$  and  $Y$  with  $X \cap Y = \emptyset$

$X$  *relevant* input variables

$Y$  *irrelevant* input variables

We are interested in the number of models projected onto  $X$ :

$$\#\exists Y.F(X, Y)$$

Example  $F(X, Y) = x \vee y$

$$X = \{x\} \quad Y = \{y\}$$

$$\mathcal{M}(\exists Y.F(X, Y)) = \{x, \neg x\}$$

$$\#\exists Y.F(X, Y) = 2$$

$$X = \{x, y\} \quad Y = \emptyset$$

$$\mathcal{M}(\exists Y.F(X, Y)) = \{xy, x\neg y, \neg xy\}$$

$$\#\exists Y.F(X, Y) = 3 = \#F(X, Y)$$



# Our Dual Approach Facilitates the Detection of Partial Models

```
$ cat clause.form
p | q | r | s
$ dualiza -e -r p,r,s clause.form
ALL SATISFYING ASSIGNMENTS
s
r !s
!r !s
$ dualiza -r p,r,s clause.form
NUMBER SATISFYING ASSIGNMENTS
8
```

```
$ dualiza -r p,r,s clause.form -l | grep RULE
c LOG 1 RULE UNX 1 -4
c LOG 1 RULE UNX 2 -4
c LOG 1 RULE BNOF 1 -4
c LOG 2 RULE UNX 3 -3
c LOG 2 RULE BNOF 2 -3
c LOG 3 RULE UNY 1 -2
c LOG 3 RULE ENO 1
```

# Dual Representation of $F(X, Y)$

$$P(X, Y)$$

|||

$$F(X, Y)$$

$$N(X, Y)$$

|||

$$\neg F(X, Y)$$

# Dual Representation of $F(X, Y)$

$$\exists S.P(X, Y, S)$$

|||

$$F(X, Y)$$

$$\exists T.N(X, Y, T)$$

|||

$$\neg F(X, Y)$$

# The General Case – Duality with Projection onto Relevant Input Variables

$$\exists Y, S. P(X, Y, S)$$

|||

$$\exists Y. F(X, Y)$$

$$\exists Y, T. N(X, Y, T)$$

|||

$$\exists Y. \neg F(X, Y)$$

# A First Example

$$F(X, Y) = (p \vee q \vee r \vee s)$$

$$X = \{p, r, s\}$$

$$Y = \{q\}$$

$$P(X, Y, S) = (p \vee q \vee r \vee s)$$

$$S = \emptyset$$

$$N(X, Y, T) = (\neg p) \wedge (\neg q) \wedge (\neg r) \wedge (\neg s)$$

$$T = \emptyset$$

# A First Example

$$F(X, Y) = (p \vee q \vee r \vee s)$$

$$X = \{p, r, s\}$$

$$Y = \{q\}$$

$$P(X, Y, S) = (p \vee q \vee r \vee s)$$

$$S = \emptyset$$

$$N(X, Y, T) = (\neg p) \wedge (\neg q) \wedge (\neg r) \wedge (\neg s)$$

$$T = \emptyset$$

Step	Rule	$I$	$P _I$	$N _I$	$M$	Found
0		()	$(p \vee q \vee r \vee s)$	$(\neg p) \wedge (\neg q) \wedge (\neg r) \wedge (\neg s)$	0	
1	UNXY	$s$	$\emptyset$	$(\neg p) \wedge (\neg q) \wedge (\neg r) \wedge ()$	0	
2	BN0F	$\neg s$	$(p \vee q \vee r)$	$(\neg p) \wedge (\neg q) \wedge (\neg r)$	4	$s$
3	UNXY	$\neg sr$	$\emptyset$	$(\neg p) \wedge (\neg q) \wedge ()$	4	
4	BN0F	$\neg s \neg r$	$(p \vee q)$	$(\neg p) \wedge (\neg q)$	6	$\neg sr$
5	UNXY	$\neg s \neg rq$	$\emptyset$	$(\neg p) \wedge ()$	6	
6	EN0	$\neg s \neg rq$	$\emptyset$	$(\neg p) \wedge ()$	8	$\neg s \neg r$

## With the Non-Dual Approach Only Total Models Are Detected

$$F(X, Y) = (p \vee q \vee r \vee s)$$

$$P(X, Y, S) = (p \vee q \vee r \vee s)$$

$$X = \{p, r, s\}$$

$$S = \emptyset$$

$$Y = \{q\}$$

# With the Non-Dual Approach Only Total Models Are Detected

$$F(X, Y) = (p \vee q \vee r \vee s)$$

$$P(X, Y, S) = (p \vee q \vee r \vee s)$$

$$X = \{p, r, s\}$$

$$Y = \{q\}$$

$$S = \emptyset$$

Step	Rule	$I$	$P _I$	$M$	Found
0		$()$	$(p \vee q \vee r \vee s)$	0	
1	DX	$s$	$\emptyset$	0	
2	DX	$sr$	$\emptyset$	0	
3	DX	$srp$	$\emptyset$	0	
4	DYS	$srpq$	$\emptyset$	0	
5	BP1F	$sr \neg p$	$\emptyset$	1	$srp$
6	DYS	$sr \neg pq$	$\emptyset$	1	
7	BP1F	$s \neg r$	$\emptyset$	2	$sr\bar{p}$
8	DX	$s \neg rp$	$\emptyset$	2	
9	DYS	$s \neg rpq$	$\emptyset$	2	
10	BP1F	$s \neg r \neg p$	$\emptyset$	3	$s \neg rp$
11	DYS	$s \neg r \neg pq$	$\emptyset$	3	
12	BP1F	$\neg s$	$(p \vee q \vee r)$	4	$s \neg r \neg p$
$\vdots$					



# Can We Compete with State-Of-The-Art #SAT Solvers?

```
$ cat clause4.form  
(x1 | x2 | x3 | x4)
```

# Can We Compete with State-Of-The-Art #SAT Solvers?

```
$ cat clause4.form
(x1 | x2 | x3 | x4)
```

Formula	Mode	sharpSAT [s]	DUALIZA [s]
clause10	dual	$< 1 \cdot 10^{-2}$	$< 1 \cdot 10^{-2}$
	block	$< 1 \cdot 10^{-2}$	$2 \cdot 10^{-2}$
	flip	$< 1 \cdot 10^{-2}$	$< 1 \cdot 10^{-2}$
clause20	block	$1 \cdot 10^{-2}$	$9 \cdot 10^{-1}$
	flip	$1 \cdot 10^{-2}$	$2 \cdot 10^{-1}$
clause30	block	$1 \cdot 10^{-2}$	$4 \cdot 10^4$
	flip	$1 \cdot 10^{-2}$	$2 \cdot 10^2$
clause100	dual	$< 1 \cdot 10^{-2}$	$< 1 \cdot 10^{-2}$
clause1000	dual	$8 \cdot 10^{-2}$	$2 \cdot 10^{-2}$
clause10000	dual	$1 \cdot 10^1$	$2 \cdot 10^{-1}$

# Where Our Dual Approach Really Wins

```
$ cat nrp4.form
(x1 | x2 | x3 | x4) |
(x5 = x2 ^ x3 ^ x4) |
(x6 = x1 ^ x3 ^ x4) |
(x7 = x1 ^ x2 ^ x4) |
(x8 = x1 ^ x2 ^ x3)
```

# Where Our Dual Approach Really Wins

```
$ cat nrp4.form
(x1 | x2 | x3 | x4) |
(x5 = x2 ^ x3 ^ x4) |
(x6 = x1 ^ x3 ^ x4) |
(x7 = x1 ^ x2 ^ x4) |
(x8 = x1 ^ x2 ^ x3)
```

Formula	Method	sharpSAT [s]	DUALIZA [s]
nrp10	dual	$9 \cdot 10^{-2}$	$< 1 \cdot 10^{-2}$
nrp20	dual	$7 \cdot 10^2$	$1 \cdot 10^{-2}$
nrp21	dual	$2 \cdot 10^3$	$1 \cdot 10^{-2}$
nrp22	dual	*	$1 \cdot 10^{-2}$
nrp100	dual	*	$8 \cdot 10^{-2}$
nrp1000	dual	*	$1 \cdot 10^1$
nrp5000	dual	*	$2 \cdot 10^2$

# Calculus

$$\text{EP0: } (P, N, I, M) \rightsquigarrow_{\text{EP0}} M \text{ if } \emptyset \in P|_I \text{ and } \text{decs}(I) = \emptyset$$

$$\text{EP1: } (P, N, I, M) \rightsquigarrow_{\text{EP1}} M + 2^{|X-I|} \text{ if } P|_I = \emptyset \text{ and } \text{var}(\text{decs}(I)) \cap X = \emptyset$$

$$\text{EN0: } (P, N, I, M) \rightsquigarrow_{\text{EN0}} M + 2^{|X-I|} \text{ if } \emptyset \in N|_I \text{ and } \text{var}(\text{decs}(I)) \cap X = \emptyset$$

$$\text{BP0F: } (P, N, I\ell^d I', M) \rightsquigarrow_{\text{BP0F}} (P, N, I\bar{\ell}^{f(m')}, M) \text{ if } \emptyset \in P|_{I\ell I'} \text{ and } \text{var}(\text{decs}(I')) = \emptyset \text{ and } m' = \sum \{m \mid \ell^{f(m)} \in I'\}$$

$$\text{JP0: } (P, N, II', M) \rightsquigarrow_{\text{JP0}} (P \wedge C^r, N, I\ell', M - m') \text{ if } \emptyset \in P|_{II'} \text{ and } P \models C \text{ and } C|_I = \{\ell'\} \text{ and } m' = \sum \{m \mid \ell^{f(m)} \in I'\}$$

$$\text{BP1F: } (P, N, I\ell^d I', M) \rightsquigarrow_{\text{BP1F}} (P, N, I\bar{\ell}^{f(m'+m'')}, M + m'') \text{ if } P|_{I\ell I'} = \emptyset \text{ and } \text{var}(\ell) \in X \text{ and } \text{var}(\text{decs}(I')) \cap X = \emptyset \text{ and } m' = \sum \{m \mid \ell^{f(m)} \in I'\} \text{ and } m'' = 2^{|X-I\ell I'|}$$

$$\text{BP1L: } (P, N, I\ell^d I', M) \rightsquigarrow_{\text{BP1L}} (P \wedge D, N, I\bar{\ell}, M + m'') \text{ if } P|_{I\ell I'} = \emptyset \text{ and } \text{var}(\ell) \in X \text{ and } \text{var}(\text{decs}(I')) \cap X = \emptyset \text{ and } m'' = 2^{|X-I\ell I'|} \text{ and } D = \pi(\neg \text{decs}(I\ell), X)$$

# Calculus

BN0F:  $(P, N, I\ell^d I', M) \rightsquigarrow_{\text{BN0F}} (P, N, I\bar{\ell}^{f(m'+m'')}, M + m'')$  if  $\emptyset \in N|_{I\ell I'}$  and  $\text{var}(\ell) \in X$  and  $\text{var}(\text{decs}(I')) \cap X = \emptyset$  and  $m' = \sum \{m \mid \ell^{f(m)} \in I'\}$  and  $m'' = 2^{|X - I\ell I'|}$

BN0L:  $(P, N, I\ell^d I', M) \rightsquigarrow_{\text{BN0L}} (P \wedge D, N, I\bar{\ell}, M + m'')$  if  $\emptyset \in N|_{I\ell I'}$  and  $\text{var}(\ell) \in X$  and  $\text{var}(\text{decs}(I')) \cap X = \emptyset$  and  $m'' = 2^{|X - I\ell I'|}$  and  $D = \pi(\neg \text{decs}(I\ell), X)$

---

DX:  $(P, N, I, M) \rightsquigarrow_{\text{DX}} (P, N, I\ell^d, M)$  if  $\emptyset \notin (P \wedge N)|_I$  and  $\text{units}((P \wedge N)|_I) = \emptyset$  and  $\text{var}(\ell) \in X - I$

DYS:  $(P, N, I, M) \rightsquigarrow_{\text{DYS}} (P, N, I\ell^d, M)$  if  $\emptyset \notin (P \wedge N)|_I$  and  $\text{units}((P \wedge N)|_I) = \emptyset$  and  $\text{var}(\ell) \in (Y \cup S) - I$  and  $X - I = \emptyset$

---

UP:  $(P, N, I, M) \rightsquigarrow_{\text{UP}} (P, N, I\ell, M)$  if  $\{\ell\} \in P|_I$

UNXY:  $(P, N, I, M) \rightsquigarrow_{\text{UNXY}} (P, N, I\bar{\ell}^d, M)$  if  $\{\ell\} \in N|_I$  and  $\text{var}(\ell) \in X \cup Y$  and  $\emptyset \notin P|_I$  and  $\text{units}(P|_I) = \emptyset$

UNT:  $(P, N, I, M) \rightsquigarrow_{\text{UNT}} (P, N, I\ell, M)$  if  $\{\ell\} \in N|_I$  and  $\text{var}(\ell) \in T$  and  $\emptyset \notin P|_I$  and  $\text{units}(P|_I) = \emptyset$

---

FP:  $(P \wedge C^r, N, I, M) \rightsquigarrow_{\text{FP}} (P, N, I, M)$  if  $\emptyset \notin P|_I$

# Conclusion and Future Work

We are on the right track

- ▶ DUALIZA is competitive on some CNF formulae and
- ▶ outperforms state-of-the-art #SAT solvers on another class of formulae.

# Conclusion and Future Work

We are on the right track

- ▶ DUALIZA is competitive on some CNF formulae and
- ▶ outperforms state-of-the-art #SAT solvers on another class of formulae.

In the near future, we plan to

- ▶ incorporate dual conflict analysis with backjumping and redundant clause learning,
- ▶ drop decision restrictions,
- ▶ capture component reasoning and
- ▶ weighted projected model counting for Bayesian reasoning,
- ▶ optimize circuit representation to improve CNF encoding, and
- ▶ explore dual preprocessing techniques.